

# Reply to ‘Comment on ‘Gravitating Magnetic Monopole in the Global Monopole Spacetime’ ’

E. R. Bezerra de Mello \*

*Departamento de Física-CCEN Universidade Federal da Paraíba 58.059-970, J. Pessoa, PB C. Postal 5.008, Brazil*

In this Reply I present some arguments in favor of the stability of the topological defect composed by global and magnetic monopoles.

In a recent paper A. Achúcarro and J. Urrestilla [1] have pointed the stability problem for composite, global and magnetic monopoles defect, presented in a previous publication [2]. In the former it is claimed that this topological defect is not classically stable against axially symmetric angular deformation on the boson field associated with the global sector <sup>1</sup>. By this deformation an extra tension is created at the north-pole which drags the core of the global monopole upwards *with no cost of energy*. Once the cores of both monopoles are separated, they are repelled by an induced self-interaction consequence of the distortion on the magnetic fields due to the solid angle deficit  $\delta\Omega = 4\pi\Delta$ , being  $\Delta$  the parameter associated with the energy scale where the gauge symmetry is spontaneously broken. However, in the composite defect system this analysis must take into account the induced self-energy more carefully. In order to infer the behavior of this self-energy I briefly comment previous analysis for the ideal case. In [5] it was calculated the formal expression to the electrostatic self-energy associated with a test charged particle in the pointlike global monopole spacetime. It was observed that it is positive, and by numerical evaluation can be seen that the induced electrostatic strength increases for larger values of the parameter  $\Delta$ . <sup>2</sup> Considering the effect of the core of the global monopole, the calculation of the self-energy becomes much more complicated. Although the exact expression for this self-energy becomes impossible to be obtained, we can infer its behavior as following: at the global monopole center,  $r = 0$ , the metric tensor is Minkowski. Its component  $g_{11}$  changes from 1 to  $1/(1 - \Delta)$  for distance  $r \geq r_{gm}$ , increasing the value of an effective solid angle deficit. On the other hand, the distortion on the electric field decreases for larger separation. So for the case where increasing the distance the augment of the solid angle deficit becomes dominant, the distortion increases with the separation up to  $r_{gm}$ ; conse-

quently we can infer that this self-energy goes to zero for very small separation between the charged particle and the core of the monopole, peaks around the distance of the order of the monopole’s size,  $r_{gm}$ , and decreases approximately with  $1/r$  for larger value of separation. Assuming that this effect is presented to magnetic charged particle as the authors also admit, to separate the cores of the global and magnetic monopoles needs some amount of energy to overcome this barrier <sup>3</sup>. So the question about the classical stability condition of the composite defect against axially symmetric perturbation must take into account this fact.

Another point raised by Achúcarro and Urrestilla is about the profile of the Higgs field. In fact for the model analysed in [2] there are three mass parameters to scale the radial distance. In our numerical analysis we adopted the mass of the boson vector:  $x = e\eta r$ . This choice of mass scale seemed convenient for us to analyse the behavior of the fields varying the self-coupling  $\lambda$ , keeping the electric coupling  $e$  fixed.

## Acknowledgment

This work was partially supported by CNPq.

- 
- [1] A. Achúcarro and J. Urrestilla, pre-print, hep-th/0212148.
  - [2] J. Spinelly, U. de Freitas and E. R. Bezerra de Mello, Phys. Rev. D **66**, 024018 (2002).
  - [3] A. S. Goldhaber, Phys. Rev. Lett. **63**, 2158 (1989).
  - [4] A. Achúcarro and J. Urrestilla, Phys. Rev. Lett. **85**, 3091 (2000).
  - [5] E. R. Bezerra de Mello and C. Furtado, Phys. Rev. D **56**, 1345 (1997).
  - [6] M. Barriola and A. Vilenkin, Phys. Rev. Lett. **63**, 341 (1989).

---

\*E-mail: emello@fisica.ufpb.br

<sup>1</sup>The stability problem of the global monopole against angular perturbation has been observed by A. S. Goldhaber [3], and also by the authors in [4].

<sup>2</sup>Although in [1] the authors said that the effect of the core has been considered, this is not true. In [5] the metric tensor considered was associated with the global monopole with no internal structure.

---

<sup>3</sup>An estimation for the peak of the magnetostatic self-energy can be provided considering the expression for self-energy given in [5] and that the monopole’s size is of order  $\eta^{-1}\lambda^{-1/2}$  [6].